




Bilkent University

Quiz # 04  
Math 101-Section 04 Calculus I  
19 October 2023 Thursday  
Instructor: Ali Sinan Sertöz  
**Solution Key**

**Q-1)**

The sides of the triangle on the right are changing as differentiable functions of time. At a particular time, say at  $t = t_0$ , we observe that  $b(t_0) = 15\text{cm}$ ,  $c(t_0) = 7\text{cm}$  and  $\theta(t_0) = \pi/3$ . We also observe that at that moment side  $a$  is decreasing at a rate of  $2\text{cm/s}$ , side  $b$  is increasing at a rate of  $1\text{cm/s}$  and side  $c$  is increasing at a rate of  $3\text{cm/s}$ . Find how fast  $\theta$  is changing at that moment.



Hint: You may find it useful to recall the cosine rule  $a^2 = b^2 + c^2 - 2bc \cos \theta$ .

Grading: 10 points

**Solution:**

From the cosine rule we find that

$$a(t)^2 = b(t)^2 + c(t)^2 - 2b(t)c(t) \cos \theta(t), . \quad (*)$$

Substituting in the values  $b(t_0) = 15\text{cm}$ ,  $c(t_0) = 7\text{cm}$  and  $\theta(t_0) = \pi/3$  we find that

$$a(t_0) = 13\text{cm}$$

Taking derivatives of both sides of (\*) with respect to  $t$  we find

$$2a(t_0)a'(t_0) = 2b(t_0)b'(t_0) + 2c(t_0)c'(t_0) - 2b'(t_0)c(t_0) \cos \theta(t_0) - b(t_0)c'(t_0) \cos \theta(t_0) + 2b(t_0)c(t_0) \sin \theta(t_0)\theta'(t_0).$$

Again putting in the given data  $a(t_0) = 13\text{cm}$ ,  $b(t_0) = 15\text{cm}$ ,  $c(t_0) = 7\text{cm}$ ,  $\theta(t_0) = \pi/3$ ,  $a'(t_0) = -2\text{cm/s}$ ,  $b'(t_0) = 1\text{cm/s}$ ,  $c'(t_0) = 3\text{cm/s}$  and recalling that  $\cos \pi/3 = 1/2$  and  $\sin \pi/3 = \sqrt{3}/2$ , we find that

$$\theta'(t_0) = -\frac{8\sqrt{3}}{35}\text{cm/s}.$$

Hence  $\theta$  is **decreasing** at the rate of  $\frac{8\sqrt{3}}{35}\text{cm/s}$  at that moment. (This is approximately  $4\text{mm/s}$ .)